

Some algebraic properties of
compact topological groups

Compact topological groups: examples

connected:

- S^1 , circle group.
- $\text{SO}(3, \mathbb{R})$, rotation group

not connected:

- Every finite group, with the discrete topology.
- $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$: inverse limit of finite Galois groups,
- \mathbb{Z}_p : inverse limit of finite cyclic groups

Such inverse limits inherit a topology from the discrete finite groups. It is

- *compact* (Tychonoff's Theorem) and
- *totally disconnected*.

Compact and tot. disconn. topological group = *profinite group* = inverse limit of finite groups

Familiar examples:

- infinite Galois groups
- matrix groups such as $\text{GL}_n(\mathbb{Z}_p)$
- free profinite groups

Theorem 1 *Let G be a compact group with identity component G^0 .*

(i) G/G^0 is a profinite group

(ii) $G^0 = Z \cdot P$ where Z is the centre of G^0 and

$$P \cong \frac{\prod S_i}{D}$$

is a Cartesian product of compact connected simple Lie groups S_i modulo a central subgroup D .

Part (ii) : Hilbert's 5th problem in compact case

Z is essentially a product of copies of S^1 : a 'protorus'

The compact connected simple Lie groups are well known – $\text{SO}(n)$, $\text{SU}(n)$ etc.

All results are joint work with **Nikolay Nikolov**.

G will denote a compact group

N is a normal subgroup of (the underlying *abstract group*) G .

Definition G is of *f.g. type* if the maximal profinite quotient G/G^0 is topologically finitely generated;

[equivalently: G/G^0 is an inverse limit of finite d -generator groups for some fixed number d .]

Theorem 2 ('Serre's question') *If G is of f.g. type and G/N is finite then N is open in G .*

Since the topology on a *profinite* group is defined by the family of all open subgroups (not true for connected groups!), an immediate consequence is

Corollary 1 (‘rigidity’) *If G is a finitely generated profinite group then every group homomorphism from G to any profinite group is continuous.*

In particular this shows that the **topology** on such a profinite group is uniquely determined by the **group-theoretic** structure. In fact it’s **definable**: in contrast to abstract groups, a f. g. profinite group is determined up to isomorphism by its *first-order theory* (Lubotzky / Jarden).

Remarks. (i) In any compact group, open subgroups all have finite index (immediate from the definition).

(ii) A compact *connected* group has *no* proper subgroups of finite index: *not* obvious from the definition but follows from the structure theory, which implies that such a group is *divisible*, i.e. all elements have *n*th roots for all *n*. So the meat of Theorem 2 is in the *profinite* case.

(iii) The restriction to *f.g. type* is absolutely necessary: in infinitely generated profinite groups the topology is only loosely connected to the abstract group structure.

Examples: here C_q is a cyclic group of order q , and p is a prime.

- The profinite group $C_p^{\mathbb{Z}}$ has $2^{2^{\aleph_0}}$ subgroups of index p , but only countably many open subgroups.

This group therefore has many distinct topologies, but the resulting topological groups are all isomorphic.

- The profinite groups $A = \prod_{n \in \mathbb{N}} C_{p^n}$ and $A \times \mathbb{Z}_p$ are isomorphic as abstract groups, but not as topological groups.

However, every finite (abstract) image of A occurs also as a continuous image.

- There is a profinite group having no abelian continuous image, but having C_2 as an abstract image.

What about **countable images**?

A compact group can't be countably infinite.

Could there be a countably infinite *abstract* image? YES!

Let A be an infinite f.g. abelian profinite group. Then either A maps onto \mathbb{Z}_p or A maps onto $B = \prod_{p \in P} C_p$, P an infinite set of primes.

We have additive group homomorphisms

$$\begin{aligned} \mathbb{Z}_p &\hookrightarrow \mathbb{Q}_p \rightarrow \mathbb{Q}, \\ B &\cong \prod_{p \in P} \mathbb{F}_p \twoheadrightarrow \prod_{p \in P} \mathbb{F}_p / \sim = F \twoheadrightarrow \mathbb{Q}, \end{aligned}$$

where F is a non-principal ultraproduct, hence a field of characteristic 0.

In both cases these compose to give a group epimorphism from A onto \mathbb{Q} .

Main result:

Theorem 3 *If G is of f.g. type and G/N is countably infinite then G/N has an infinite virtually-abelian quotient.*

This implies for example that G cannot map onto a *countably infinite simple group*.

Corollary 2 *Suppose G is of f.g. type. Then G has a countably infinite abstract image if and only if G has an infinite virtually-abelian continuous quotient.*

By Theorem 2, if G/N is residually finite then N is closed. As every finitely generated abelian group is residually finite, we get

Corollary 3 *If G/N is finitely generated (as abstract group) then G/N is finite (and so N is open in G).*

A generalisation

Let \overline{N} denote the closure of N in G .

$$\begin{aligned} G/N \text{ countable} &\implies G/\overline{N} \text{ a countable compact group} \\ &\implies G/\overline{N} \text{ finite} \implies \overline{N} \text{ open.} \end{aligned}$$

Definition

N is *virtually dense* in G if \overline{N} is open in G .

N can have infinite index – i.e. $N < \overline{N}$ – if G is *abelian*.

Another example: I an infinite index set,

$$G = \prod_{i \in I} H_i$$

a product of non-trivial compact groups. Then the *restricted direct product*

$$N = \bigoplus_{i \in I} H_i$$

is dense in G , and has infinite index.

Definition

G as above is *strictly infinite semisimple* if I is infinite and each of the H_i is either a *finite simple group* or a *connected simple Lie group*.

Theorem 4 *Let G be a compact group of f.g. type. Then G has a virtually-dense normal subgroup of infinite index if and only if G has a (continuous) quotient that is either*

- *infinite and virtually abelian or*
- *virtually (strictly infinite semisimple).*

We can also characterize precisely those G that have a *proper dense* normal subgroup: the answer involves certain restrictions on the simple factors occurring in the strictly infinite semisimple quotient.

Exercise. Deduce: if G is just-infinite and not virtually abelian then *every normal subgroup of G is closed* (and so G is just-infinite as abstract group).

Key question: how to get *topological* information from *algebraic* input?

Given: (i) definition of **topological group**:
group multiplication is continuous,

(ii) the definition of **compact**, which implies that *a continuous image of a compact set is compact, hence closed*.

Lemma 1 *Let X be a closed subset of a compact group G , with $1 \in X = X^{-1}$. Then the subgroup $\langle X \rangle$ generated (algebraically) by X is closed in G if and only if there exists n such that*

$$\begin{aligned} \langle X \rangle &= X^{*n} \\ &= \{x_1 \dots x_n \mid x_i \in X\}. \end{aligned}$$

In this case, we say that X has *width* (at most) n in G , and write

$$m_X(G) \leq n.$$

If G is *profinite*,

$$m_X(G) = \sup \{m_{XK/K}(G/K) \mid K \triangleleft_o G\}.$$

– Reduces the study of $m_X(G)$ to the case where G is **finite**.

Results on finite groups

A finite group Q is *almost-simple* if

$$S \leq Q \leq \text{Aut}(S)$$

for some simple (non-abelian) group S

Definition. For a finite or profinite group G ,

$$G_0 = \bigcap \{K \triangleleft_o G \mid G/K \text{ is almost-simple}\}.$$

Theorem 5 $H \triangleleft G$, a finite group.

Y a symmetric subset of G such that

$$H \langle Y \rangle = G' \langle Y \rangle = G.$$

If $H \leq G_0$ or $\langle Y \rangle = G$ then

$$[H, G] = \left(\prod_{y \in Y} [H, y] \right)^{*f}$$

where $f = f(d, r) = O(r^6 d^6)$, $d = d(G)$,

Here $[H, G] = \langle [h, g] \mid h \in H, g \in G \rangle$,

$G' = [G, G]$, derived group of G).

Routine compactness arguments turn Theorem 5 into

Theorem 6 *G f. g. profinite, $H \triangleleft G$ closed. Y a finite symmetric subset of G such that*

$$\overline{H \langle Y \rangle} = \overline{G' \langle Y \rangle} = G.$$

If $H \leq G_0$ or $\overline{\langle Y \rangle} = G$ then

$$[H, G] = \left(\prod_{y \in Y} [H, y] \right)^{*f}$$

for some finite f .

Corollary 4 *G f. g. profinite, $H \triangleleft G$ closed. Then $[H, G]$ is closed.*

Also true (but harder) when G is compact of f.g. type.

Suppose that $Y \subseteq N \triangleleft G$. Then $[H, G] \leq N$. It is now easy to deduce the key ‘**reduction theorem**’:

Corollary 5 *Let G be a finitely generated profinite group with a normal subgroup N . If*

$$NG' = NG_0 = G$$

then $N = G$.

Reduces problems about G to problems about

- G/G' : an **abelian** group,
- G/G_0 : an extension of a **semisimple** group by a **soluble** group. In fact there are closed normal subgroups

$$A_0 \triangleleft A_1 \triangleleft A_2 \triangleleft A_3 = G/G_0$$

with A_0 semisimple, A_i/A_{i-1} abelian ($i = 1, 2, 3$)

Typical applications:

(1) Quick proof of Theorem 2 (‘Serre’s question’).

The abelian case is easy, and the semisimple case follows from

Theorem (Martinez/Zelmanov, Saxl/Wilson, 1996-97) *Let $q \in \mathbb{N}$. In any finite simple group S , the set $\{x^q \mid x \in S\}$ has width at most $f(q)$, a finite number depending only on q .*

(2):

Theorem *G f.g. profinite, $q \in \mathbb{N}$. Then G^q is open in G .*

(3): Proofs of Theorems 3 and 4: reduction to problems about semisimple groups.

Normal subgroups in semisimple groups.

This depends on some different ideas. Suppose

$$G = \prod_{i \in I} S_i$$

where I is an infinite index set and each S_i is a finite simple group, appearing with finite multiplicity. To each non-principal ultrafilter \mathcal{U} on I we associate a certain normal subgroup $K_{\mathcal{U}}$ of G , and prove:

$$|G/K_{\mathcal{U}}| \geq 2^{\aleph_0}.$$

($G/K_{\mathcal{U}}$ is what is known as a ‘metric ultraproduct’; it is a simple group.)

Proposition 1 *Let N be a proper normal subgroup of G . If N is dense in G then $N \leq K_{\mathcal{U}}$ for some non-principal ultrafilter \mathcal{U} .*

Together with a similar construction in the case where the S_i are simple compact Lie groups, it gives

Theorem 7 *Let G be a semisimple compact group of f.g. type and N a normal subgroup of G . If $|G/N|$ is infinite then $|G/N| \geq 2^{\aleph_0}$.*

N. Nikolov, D. Segal, ‘Generators and commutators in finite groups; abstract quotients of compact groups’, *Invent. Math.* (2012)

N. Nikolov, D. Segal, ‘On normal subgroups of compact groups’, *JEMS*, to appear